

- With **Bromadiolone**
- Single Feed Rodenticide
- Palatable and Attractive

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE Contains Denatonium Benzoate (Bittering Agent)

# FOR CONTROL OF RATS & MICE

PRIMARY PACK CONTAINS 80 x 10 g SACHETS WHICH ARE ILLEGAL TO SELL SEPARATELY. DO NOT DESTROY CONTAINER WHILE PRODUCT STILL REMAINS.

Net 800 g





TOMCAT is a trademark of Bell Laboratories, Inc., U.S.A.



Australian Agent: Barmac

Barmac
A Division of Amgrow Australia Pty Ltd
(ABN 22 069 900 456)
82 Christensen Road, Stapylton, QLD 4207
EMERGENCY CONTACT NO. 1800 AMGROW (1800 264 789) | www.barmac.com.au APVMA APPROVAL NO: 86331/120939

RESTRAINTS: DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.

DO NOT place bait sachets in the open.

DO NOT use bait stations further than 2 m from buildings.

### DIRECTIONS FOR USE

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, agricultural, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around rain terminals	Rats (Rattus norvegicus, Rattus rattus) Mice (Mus musculus)	2 to 20 x 10 g soft baits per bait station or placement. 1 to 4 x 10 g soft baits per bait station or placement.	Eliminate as far as practiceable, all alternative food sources Bail areas where conditions might encourage rodent activity. Place bails so as to prevent access to non-target animals Tamper resistant bail stations must be used at locations where access to non-target animals cannot otherwise be prevented. Inspect bail stations on a regular basis. Replace eaten or old/deteriorated bails. If eaten quickly, increase the number of bail stations. Confirme observation and replenish until no more bails are taken. Baiting for at least two week is necessary to reduce ratifrouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoilt bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead animals and rodent faeces by rat/mouse numbers. burning or burial, and left over balt as per disposal instructions. Baiting for at least two weeks is necessary to reduce

### PRECAUTIONS

# DO NOT remove bait from sachet.

The use or union to the second of the second The use of this product for the control of protected native rodents requires permission from wildlife authorities. DO NOT place bait in places accessible to children, domestic animals livestock and non-target wildlife, unless in



# FOR CONTROL OF RATS & MICE

TOMCAT Soft Bait is a highly palatable soft bait for the control of rats and mice including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

Ready to use bait - Kills in a single feed. Soft Bait = palatable and attractive.

## PRECAUTIONS continued from side panel.

DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed, or surfaces that come into direct contact with food. The product is restricted to use in an around buildings (within 2 metres) or enclosed spaces.

DO NOT apply bait to crops.

### PROTECTIONS

PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT DO NOT contaminate dams, rivers, streams or drains with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product and dead or affected rodents and rodent faeces are hazardous to pigs,

### cats, dogs, native wildlife and poultry. PROTECTION OF LIVESTOCK

Bait, and dead or affected rodents are hazardous to birds, dogs, cats, pigs, poultry and non-target wildlife. Regularly search for, remove and destroy dead or affected rodents.

# STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area, as cool as possible.

DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away form children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. To the extent possible, untaken baits should be destroyed according to State or Territory requirements. To the extent possible, animal carcasses should be recovered and destroyed according to State or Territory requirements.

Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear rubber gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K1 (Phytomenadione) is antidotal.

# FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

For further information please refer to the Safety Data Sheet which can be obtained from the supplier.

In a transport emergency dial 000, Police or Fire Brigade. For specialist advice in an emergency only, call (02) 9037 2994 (24 h). CONDITIONS OF SALE

The use of TOMCAT® Soft Bait is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose, Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Competition and Consumer Act 2010.

# **GHS Hazard and Precautionary Statements**

May cause damage to organs through prolonged or repeated exposure. Toxic to aquatic life with long lasting effects.

Batch No.:

Do not get in eyes, on skin, or on clothing. Do not eat, drink or smoke when using this product. Avoid release to the environment. Use personal protective equipment as required. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Collect spillage. Not combustible. Use extinguishing media suited to burning materials. Dispose of small quantities and empty containers by wrapping with paper and putting in garbage. For larger quantities, if recycling or reclaiming is not possible, use a commercial waste disposal service.

D.O.M.:

POISON
KEEP OUT OF REACH OF CHILDREN
TOMCAT® SOFT BAIT
ACTIVE CONSTITUENT: 0.05 glvg BROMADIOLONE
Contains Denatonium Berzoate (Bittering Agent)
Before use read all directions on the outer pack.
APVMA Approval No.: 86331/120939 NET 10 g

NOT TO BE SOLD SEPARATELY

2612